

Single Particle Absorption Measurements in the Mid-Infrared by Exploiting Elastic Scattering

Kevin Aptowicz, Yong-Le Pan, and Richard K. Chang
Yale University

Ronald G. Pinnick, Steven C. Hill, Kris Gurton, and Richard L. Tober
Army Research Labs

Burt V. Bronk
AFRL at US Army SBCCOM

Joint Service Scientific Conference on Chemical & Biological Defense Research


Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 01 OCT 2005		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Single Particle Absorption Measurements in the Mid-Infrared by Exploiting Elastic Scattering				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Yale University				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM001851, Proceedings of the 2003 Joint Service Scientific Conference on Chemical & Biological Defense Research, 17-20 November 2003. , The original document contains color images.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 13	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Goal: Measure the infrared absorption and scattering cross-sections for single biological and chemical aerosol particles.

Use: Single particle measurements are necessary for detailed modeling and understanding of test results from infrared stand-off detection systems.

Technique:

Two-dimensional Angular Optical Scattering (TAOS)



TAOS patterns depend upon particle shape, size, and **complex refractive index**

Coordinates for TAOS patterns

Scattering Parameters



Diameter: 54.2 μm

Refractive Index: $1.342 + i * 0.00$






Wavelength: 3.41 μm

Size Parameter: $2\pi * a / \lambda \cong 50$

Laser Polarization: Vertical




Extrapolate absorption cross-sections of spherical particles by comparison with Mie theory



Changes in absorption leads to changes in the scattering profile beyond just a scale factor.




Experimental set-up to collect TAOS patterns of droplets



Collected TAOS patterns of droplets




H_2O

Droplet Diameter: $57.4\ \mu\text{m}$
Refractive Index: $1.405 + i\ 0.018$






50% H_2O - 50% D_2O

Droplet Diameter: $54.2\ \mu\text{m}$
Refractive Index: $1.342 + i\ 0.010$



D_2O


Droplet Diameter: $55.2\ \mu\text{m}$
Refractive Index: $1.279 + i\ 0.002$



Comparison between experiment and Mie theory


D_2O

Droplet Diameter: $55.2\ \mu m$
Refractive Index: $1.279 + i\ 0.002$







50% H_2O - 50% D_2O

Droplet Diameter: $54.2\ \mu m$
Refractive Index: $1.342 + i\ 0.010$



H_2O


Droplet Diameter: $57.4\ \mu m$
Refractive Index: $1.405 + i\ 0.018$



Current work


Collaboration with MIT Lincoln Lab

(Anish Goyal, Tom Jeys, and Antonio Sanchez)







Large Angle Two-dimensional Angular Optical Scattering

LA TAOS patterns collected
in the **visible** of clusters
($\lambda = 532$ nm)




LA TAOS patterns collected
in the **mid-IR** of clusters
($\lambda = 3.9 \mu\text{m}$)

Variability within a data set is due to multiple factors: cluster size, shape, and orientation, and optical alignment distortions.




LA TAOS in the visible and mid-infrared




By increasing the wavelength,
the LA TAOS technique
becomes more sensitive to
larger structure sizes.



SEM of Tryptophan




Visible LA TAOS pattern of Tryptophan
at $\lambda = 532$ nm



Mid-IR LA TAOS patterns of Tryptophan at $\lambda = 3.9$ μm

Future Plans: Capture Dual Wavelength LA TAOS



Use two mid-infrared wavelengths to simultaneously illuminate an aerosol, then compare the LA TAOS patterns to ascertain if there is absorption at either wavelength.

Summary of Work

- Detected TAOS patterns of single 50 μm droplets composed of H_2O , D_2O , and $\text{H}_2\text{O}/\text{D}_2\text{O}$ mixture.
- Able to achieve decent visible match with results derived from Mie theory.
- Unable to implement a minimization routine to find absorption because of aberration in the collection optics as well as an inability to determine absolute angle reference.
- Collected LA TAOS patterns of Arizona Road Dust, BG, Bovine Albumin, PSL sphere cluster, and Tryptophan